

Phylogeography of the blue land crab, *Cardisoma guanhumi* (Decapoda: Gecarcinidae) along the Brazilian coast

JOSÉ F. DE OLIVEIRA-NETO^{1,3}, MARCIO R. PIE^{1,3}, MARCELO A. CHAMMAS³, ANTONIO OSTRENSKY^{2,3}
AND WALTER A. BOEGER^{1,3}

¹Departamento de Zoologia, Universidade Federal do Paraná, Curitiba, PR, Brazil, ²Departamento de Zootecnia, Universidade Federal do Paraná, Curitiba, PR, Brazil, ³Grupo Integrado de Aqüicultura e Estudos Ambientais, Universidade Federal do Paraná, Curitiba, PR, Brazil

*Several crab species are important fishery resources in Brazil and their overexploitation has led to severe population declines in some regions. The management of populations of these species should take into account the degree of gene flow among populations in different estuaries. The goal of the present study is to assess the degree of geographical structure in the genetic diversity of the blue land crab, *Cardisoma guanhumi*, along the Brazilian coast. A fragment of the control region of the mtDNA (750 bp) was sequenced for 95 specimens collected across 5 Brazilian states. Analyses using F-statistics failed to indicate any evidence of geographical structure, a result that was corroborated by a nested clade analysis of the same dataset. Mismatch distribution analyses indicated that populations of the blue land crab have experienced an expansion during their recent evolutionary past. The obtained results are similar to those recently described for another sympatric crab, *Ucides cordatus*, particularly with respect to the extensive degree of gene flow. However, populations of *C. guanhumi* seem to be older than those of *U. cordatus* and do not show the north–south expansion found in that species.*

Keywords: population genetics, d-loop, mangrove, estuary, guaiamum

Submitted 7 November 2007; accepted 22 April 2008; first published online 22 July 2008

INTRODUCTION

The blue land crab, *Cardisoma guanhumi* (Brachyura: Gecarcinidae) is a species highly adapted to terrestrial life that is found in estuarine areas throughout tropical and subtropical regions in the Atlantic Coast of the New World, from the state of Florida (USA) to the state of Santa Catarina (southern Brazil). It is an important food resource due to its large body size, which often exceeds 11 cm in carapace length. In Brazil, its highest consumption takes place in the north-eastern states, where it has considerable economic importance (Amaral & Jablonski, 2005; Ministério do Meio Ambiente, 2005). *Cardisoma guanhumi* is not considered to be under threat of extinction, yet in some areas it became rare and its average body size decreased (Amaral & Jablonsky, 2005), resulting in a reduction in its commercialization and reducing the importance of this fishery resource. As a consequence, *C. guanhumi* has been included in the list of overexploited species of the Brazilian Ministry of Environment (Ministério do Meio Ambiente, 2004).

The life-cycle of *C. guanhumi* is presumably similar to other semi-terrestrial crustaceans that live in or near mangrove areas. During the reproductive season, such crabs release thousands of larvae into the water during high tide of full moon. These larvae are carried to open waters, where

they remain for several weeks until returning to an estuary (Gifford, 1962). Genetic flow between populations of the species is accomplished by the dispersal of larvae, since the adults present reduced horizontal displacements. The distance between mangroves, the flow of water currents, and the shape of the estuaries vary considerably along the Brazilian coast, yet the influence of these factors on the dispersal potential of planktonic larvae is poorly known.

An understanding of the degree of gene flow, genetic diversity, and effective population size is the basis for the establishment of efficient conservation strategies (Frankham *et al.*, 2002). It has been usually thought that animals with this type of life cycle, with pelagic and marine larvae, would show little genetic differentiation due to the high level of gene flow. However, studies on brachyurans have shown genetic differentiation levels ranging from negligible (McMillen-Jackson & Bert, 2004; Cassone & Boulding, 2005; Pfeiler *et al.*, 2005; Oliveira-Neto *et al.*, 2007, 2008) to intermediate or high (Weber & Levy, 2000; Fratini & Vannini, 2002; Weinberg *et al.*, 2003; Roman & Palumbi, 2004).

A recent study on the ocypodid *Ucides cordatus*, a species largely sympatric with the blue land crab, failed to show any evidence of genetic structure in populations south of the River Amazon (Oliveira-Neto *et al.*, 2007). The goal of the present study is to characterize and compare the genetic structure of *C. guanhumi* along the Brazilian coast using sequences of the control region of the mitochondrial DNA, a fragment characterized by high mutation rate that is particularly suitable for population genetic studies.

Corresponding author:
M.R. Pie
Email: pie@ufpr.br

MATERIALS AND METHODS

Samples were obtained from five states along the Brazilian coast: Rio Grande do Norte (N = 10), Pernambuco (N = 19), Sergipe (N = 21), Rio de Janeiro (N = 20), and Santa Catarina (N = 23) (Figure 1). Distances between the studied estuaries vary from 400 to 5000 km and are separated from one another by sandy beaches and other estuaries. The approximately 5000 km of coast encompassed in the present study represent approximately a third of the distribution of *C. guanhumi*. There is a bifurcation of the South Equatorial Current in north-eastern Brazil near 16°S, with one branch heading northwards as the Guyana Current and the other heading south as the Brazil Current, a phenomenon that could potentially affect larval distribution, even though near-shore waters can be more influenced by coastal currents that are more variable in space and time (Miranda, 1970; Mesquita & Hatari, 1969; Silva *et al.*, 2005).

A fragment of muscle tissue of the pereiopods was removed, preserved in an EDTA–DMSO buffer (Seutin *et al.*, 1991), and maintained at –20°C. Genomic DNA was extracted using the ChargeSwitch® kit (Invitrogen) according to the manufacturer's instructions. The primers 12SUCAF3 (5'–CCA GTA NRC CTA CTA TGT TAC GAC TTA T–3') and ILEUCAR3 (5'–GCT AYC CTT TTA AAT CAG GCA C–3') were used to amplify a ≈1.6 kb fragment including the entire control region of the mtDNA (Oliveira-Neto *et al.*, 2007). Each 25-μl PCR solution had the following final concentrations: 6 mM of MgCl₂, 0.25 mM of each dNTP, 0.1 U/μl of *Taq* polymerase, 1X buffer, 2 μM of each primer, and 1.2 ng/μl of genomic DNA. Thermocycling settings included the following steps: 95°C for

2 minutes, followed by 35 cycles of 95°C for 20 seconds, 56°C for 30 seconds, and 72°C for 90 seconds, followed by a final extension of 72°C for 2 minutes. A 2 μl aliquot of each PCR product was electrophoresed in a 1.5% agarose gel, stained with EtBr, and photographed under UV-light. Successfully amplified products were purified using the MinElute kit (Qiagen) and sequenced using the internal primers ILEUCAR2: 5'–CCT TTT AAA TCA GGC ACT ATA–3', and DLUSSAF1: 5'–GTA TAA CCG CGA ATG CTG GCA C–3') (Oliveira-Neto *et al.*, 2007), generating a ≈850 pb fragment. Each 10 μl cycle-sequencing reaction included the following: 0.16 μM of primer, 0.15X of buffer, 0.5 μl of BigDye (Applied Biosystems), and 20 ng of the PCR product. The resulting solution was purified using Sephadex G50 and processed using an ABI3130 automatic sequencer. The obtained sequences were aligned unambiguously by eye.

The occurrence of geographical structure in the genetic variability of populations was tested using F-statistics and analysis of molecular variance (AMOVA). The null distribution of pairwise F_{st} values under the hypothesis of no difference between the populations was obtained by permuting haplotypes between populations. Similarly, the AMOVA was calculated based on a matrix of haplotype distances, thus allowing for comparing groups of samples. Samples from Rio de Janeiro and Santa Catarina were grouped as 'southern samples', whereas those from the remaining states were grouped as 'northern samples', to increase the statistical power of the AMOVA. This distinction took into account the distance among the collection points and their respective climatic and geomorphological characteristics, given that the regions where southern samples were obtained are colder and have fewer estuaries than the regions in the north. These analyses were complemented by a nested clade analysis (NCA), as implemented in the program GeoDis (Posada *et al.*, 2000) according to the methods of Templeton *et al.* (1995). This program calculates the clade distances D_c (average distance between the location of the members of the clade and the geographical centre of the clade) and the nested clade distances D_n (the average spatial distance between the members of each clade and the geographical centre of the entire nesting clade). Additionally, the measures of average distance between tip and interior clades within the nested group ($Int-Tip$)_c and the tip to interior distance for the nesting clade ($Int-Tip$)_n are estimated. The assessment of whether any of these distances are significantly smaller or larger than expected by chance was carried out by 10,000 permutation resamplings. Interpretation of the results followed the method given in Templeton *et al.* (1995).

Inferences on past demographic history of the blue land crab were assessed using the mismatch distribution analysis (Slatkin & Hudson, 1991; Rogers & Harpending, 1992). Three parameters are estimated using Rogers & Harpending's (1992) model: $\theta_0 = 2N_0u$, $\theta_1 = 2N_1u$, and $\tau = 2ut$, where an initial population of effective size N_0 is assumed to grow rapidly to a new size of N_1 at a time t generations before the present, and u is the per-generation probability that a mutation strikes a particular nucleotide in the region under study. These parameters were estimated using the generalized non-linear least-square approach developed by Schneider & Excoffier (1999). Also, the degree of approximation between the observed mismatch distribution and that expected under population growth was tested using Harpending's (1994) raggedness statistic.




Fig. 1. Sites included in the present study for the collection of specimens of *Cardisoma guanhumi* along the Brazilian coast. Symbols indicate the following states, from top to bottom: Rio Grande do Norte, Pernambuco, Sergipe, Rio de Janeiro and Santa Catarina. The same symbols are used in Figure 2.

two in Rio de Janeiro and two in Santa Catarina (Table 1). No haplotype was shared among locations. Nucleotide diversity was around 0.03 in all locations, with the average number of nucleotide differences ranging from 21.3 in Sergipe to 24.9 in Rio Grande do Norte.

There was no statistically significant geographical structure in the pattern of genetic variability among the studied locations according to the NCA (Figure 2; Table 2). The lowest observed *P* value was 0.035, which becomes non-significant after adjusting the critical value to account for

Table 2. Nested clade analysis of the variation in the control region of the mtDNA of *Cardisoma guanhumi* (Dc, average distance between the location of the members of the clade and geographical centre of the clade; Dn, the average spatial distance between the members of each clade and the geographical centre of the entire nesting clade. I-T, the average distance between tip and interior clades within the clade or nested group).

Clade	Dc	Dn	Clade	Dc	Dn	Clade	Dc	Dn	Clade	Dc	Dn
2-3	0	2080									
2-4	0	1270				3-2	1693	1299			
2-5	0	1549									
2-6	0	1685									
2-7	0	2065									
2-8	1940	1477	3-3	2672	1427						
2-9	0	1235				4-1	1420	1272			
2-10	0	1940	3-4	1552	1226						
2-11	0	1293	I-T	120	201						
2-12	0	1104							5-1	1393	1290
2-13	0	818									
2-14	0	1616	3-5	1169	1250						
2-15	2200	736									
I-T	-2200	-747									
2-16	0	1080				4-2	1365	1300			
2-17	0	1367									
2-18	2540	1376									
2-19	0	1445	3-6	1353	1327						
2-20	0	1168									
2-21	0	1367									
2-22	1280	1150									
I-T	-716	151	I-T	-185	-77						
2-23	0	1448									
2-24	0	1447				3-7	1546	1302			
2-25	2400	1600									
I-T	2400	-153									
2-26	0	2215									
2-27	0	1535									
2-28	1693	1597	3-8	1798	1471						
I-T	-1693	278									
2-29	0	1600									
2-30	0	1600	3-9	1600	1206						
2-31	0	1600									
2-32	0	1600	3-10	1600	1206						
I-T	0	0				4-3	1495	1341			
2-33	0	991									
2-34	1693	1507									
2-35	0	1620									
2-36	0	991	3-11	1572	1386						
2-37	0	1067							5-2	1422	1309
I-T	847	287	I-T	137	-63						
2-38	0	1500									
2-39	0	1730									
2-40	0	2770	3-12	2000	1383						
I-T	0	750									
			2-73	0	813						
2-41	0	1600									
2-42	0	800									
2-43	0	800	3-13	1280	1065						
I-T	0	-800				4-4	1266	1266			
2-44	0	2200									
2-45	0	2200	3-14	2200	1372						
I-T	0	0									
2-46	0	1940									
2-47	0	1940	3-15	1940	1189						
			3-16	0	937						

continued

Table 2. Continued

Clade	Dc	Dn															
2-48	0	600	3-17	600	1715	4-5	1500	1421	5-3	1472	1325						
2-49	0	600															
1-T	0	0															
2-50	0	3000															
2-51	0	3000															
2-52	0	1304															
2-53	340	936															
2-54	0	1576															
2-55	0	936															
2-56	0	1576															
2-57	0	533															
2-58	0	800															
2-59	0	1508										3-21	1493	1337	4-6	1495	1423
2-60	0	1388															
2-61	0	1268															
2-62	0	1268															
2-63	0	1540															
1-T	240	-138															
2-64	800	1600															
2-65	2200	1575															
2-66	0	1850															
1-T	666	67															
2-67	0	1570															
2-68	0	1270															
2-69	0	2240															
1-T	0	635															
			3-23	1693	1337												
			1-T	51	-51				1-T	-8.6	2.6						

the high number of separate statistical tests involved in NCA using a procedure such as the Bonferroni correction. Similar results were obtained using F-statistics, where none of the F_{st} values differed significantly from zero (Table 3), as well as according to the AMOVA (Table 4).

The mismatch distribution was consistent with a simple unimodal pattern, either when all samples were combined or when each location was analysed separately, as shown by the non-significant raggedness indices (Figure 3; Table 1). Such a pattern is an indication of the occurrence of population expansion. This conclusion is supported by the negative values of all estimates of Tajima's D and Fu's F_s , although only the latter reached statistical significance (the only exception is the population in Rio Grande do Norte, which might be artefactual due to the smaller sample size).

An interesting pattern becomes evident when one compares the estimates of τ along a latitudinal axis. As indicated above, τ is directly proportional to the per-generation probability of mutation in the studied DNA fragment multiplied by the time of population expansion in terms of number of generations. The oldest populations are located in the northernmost and

Table 3. F_{st} estimates between the studied samples of *Cardisoma guanhumii*. All estimates did not differ statistically from 0 ($P > 0.05$ in all comparisons). RN, Rio Grande do Norte; PE, Pernambuco; SE, Sergipe; RJ, Rio de Janeiro; SC, Santa Catarina.

	RN	PE	SE	RJ	SC
RN	0				
PE	-0.007	0			
SE	0.025	-0.006	0		
RJ	-0.004	-0.013	-0.011	0	
SC	0.016	0.002	-0.005	0.007	0

Table 4. Analysis of molecular variance (AMOVA) based on a distance matrix between haplotypes. The studied samples were separated between southern (Rio de Janeiro and Santa Catarina) and northern states (Rio Grande do Norte, Pernambuco and Sergipe).

Source of variation	DF	SS	Component variation	Percentage of variation
Among groups	1	10	-0.08	-0.55
Among populations in each group	3	42	0.002	0.01
Within populations	82	1.149	1.401	100




Fig. 3. Variation in estimates of τ among the studied sites (solid line) and their respective 95% confidence intervals.

southernmost sites, whereas the youngest population is located in the latitudinally intermediate location (Sergipe) (Figure 4). However, the broad overlap among confidence intervals suggests that this result should be interpreted with caution.




Fig. 4. Mismatch distribution analysis of the studied fragment of mtDNA of *Cardisoma guanhumii* for each location, as well for the combined dataset.

DISCUSSION

Populations of the blue land crab are characterized by a high level of genetic variability that is homogeneously distributed throughout the entire studied region along the Brazilian coast. High genetic diversity is thought to result from a combination of high mutation rate in the study fragment and the large population size of the species (Avise *et al.*, 1984), as observed in other crustacean species (e.g. McMillen-Jackson & Bert, 2004; Diniz *et al.*, 2005; Cassone & Boulding, 2006). Moreover, the unimodal mismatch distributions and the estimates of Fu's F_s indicate an inferred population expansion during the recent evolutionary history of *C. guanhumii*. The patterns observed in this species are particularly comparable to those obtained for *U. cordatus* (Brachyura: Ocypodidae) in a study carried out using the same DNA fragment (Oliveira-Neto *et al.*, 2007). These species share similar geographical ranges, from southern USA to southern Brazil, as well as several biological characteristics, such as a semiterrestrial habit near estuarine waters which are used for both spawning and larval development (Oliveira, 1946; Melo, 1996; Hill, 2001). Both species do not seem to experience obstacles for gene flow in the region encompassed by their studies. The dispersal capacity of their larvae is probably amplified through coastal currents, such that the genetic compositions of populations in different estuaries are homogenized. It is important to note that successive migration among adjacent estuaries should be sufficient for such effect, so that dispersal over very long distances in a single generation does not have to be invoked.

The implications of the extensive gene flow throughout the distribution of the blue land crab have important implications for their conservation. First, populations along the coast do not seem to be organized into discrete evolutionarily significant units (Moritz, 1994), which greatly facilitates management efforts. For instance, severely depressed populations can be enhanced by restocking efforts through laboratory mass rearing and adult translocation, so that movement of individuals among geographical regions should not compromise the genetic composition of local populations. However, gene flow over evolutionary time does not necessarily mean that populations are strongly linked in demographic terms, given that recruitment among adjacent estuaries might in

fact take place at a very low rate. Recent Bayesian methods to estimate recent migration rates might be particularly useful to elucidate this issue (Wilson & Rannala, 2003).

In spite of the above mentioned similarities, *C. guanhumii* and *U. cordatus* show intriguing differences in the inferred population dynamics. Both species show clear evidence of unimodal mismatch distributions, which are consistent with population expansion. However, the estimates of τ , the parameter that describes the time since the onset of population expansion, are markedly different between both species, with the expansion of *C. guanhumii* being considerably older than that of *U. cordatus* (21.5 and 16.6, respectively). The latitudinal variation in the estimates of τ are also markedly different, with *U. cordatus* showing a clear decrease in τ toward higher latitudes (Oliveira-Neto *et al.*, 2007), whereas the present study indicated the lowest τ estimates at intermediate latitudes. We hypothesize that the latter might represent a secondary subdivision in the range of the species, as opposed to the simple north-to-south expansion in the case of *U. cordatus*. Finally, the estimates of θ , using all samples were also markedly different (191.6 in *U. cordatus* and 514.0 in *C. guanhumii*). This parameter is proportional to the population size after the expansion multiplied by the per-generation mutation rate for the studied fragment. Given that both studies focused on the same DNA fragment and that the species are members of phylogenetically close families, there is no reason to suspect that differences in mutation rate alone would account for the obtained results. Rather, there are important ecological differences between *U. cordatus* and *C. guanhumii* that might account for the observed discrepancies. It has been known for long that *C. guanhumii* is not as strongly associated with the intertidal region as *U. cordatus*, frequently showing preference for the drier conditions of the upper shore (e.g. Oliveira, 1946 and references therein). Consequently, the population responses of each species to changes in mangrove areas would be markedly different. Further genetic studies on other crustacean species, as well as on species of mangrove trees, are necessary to discriminate species-specific features from the evolutionary dynamics of the mangrove environment itself.

ACKNOWLEDGEMENTS

The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) provided partial funding for the study and assistantships to M.R.P., W.A.B., and J.F.O.N. Additional funding was provided by the Government of the State of Paraná through the Secretaria de Estado da Ciência, Tecnologia e Ensino Superior (SETI). We are grateful to a variety of colleagues who provided samples from Rio Grande do Norte and Pernambuco (M.W. Pil), Sergipe (R.O. Ribeiro) and Santa Catarina (T.V. Oliveira and G.O. Langner). Luciana Patella provided valuable laboratory support.

REFERENCES

- Amaral A.C.Z. and Jablonski S. (2005) Conservação da biodiversidade marinha e costeira no Brasil. *Megadiversidade* 1, 43–51.

- Avise J.C., Neigel J.E. and Arnold J.** (1984) Demographic influences on mitochondrial DNA lineage survivorship in animal populations. *Journal of Molecular Evolution* 20, 99–105.
- Cassone B.J. and Boulding E.G.** (2006) Genetic structure and phylogeography of the lined shore crab, *Pachygrapsus crassipes*, along the north-eastern and western Pacific coasts. *Marine Biology* 149, 213–226.
- Diniz F.M., Maclean N., Ogawa M., Cintra I.H.A. and Bentzen P.** (2005) The hypervariable domain of the mitochondrial region in Atlantic spiny lobsters and its potential as a marker for investigating phylogeographic structuring. *Marine Biotechnology* 7, 462–473.
- Excoffier L., Laval G. and Schneider S.** (2005) Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1, 47–50.
- Frankham R., Ballou J.D. and Briscoe D.A.** (2002) *Introduction to conservation genetics*. Cambridge: Cambridge University Press.
- Fratini S. and Vannini M.** (2002) Genetic differentiation in the mud crab *Scylla serrata* (Decapoda: Portunidae) within the Indian Ocean. *Journal of Experimental Marine Biology and Ecology* 272, 103–116.
- Fu Y.X.** (1997) Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics* 147, 915–925.
- Gifford C.A.** (1962) Some observations on the general biology of the land crab, *Cardisoma guanhumi* (Latneilla), in South Florida. *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 123, 207–223.
- Harpending R.C.** (1994) Signature of ancient population growth in low-resolution mitochondrial DNA mismatch distribution. *Human Biology* 66, 591–600.
- Hill K.** (2001) 'Cardisoma guanhumi' (On-line). Accessed 28 December 2008 at http://www.sms.si.edu/IRLspec/Cardis_guanhu.htm.
- McMillen-Jackson L. and Bert T.M.** (2004) Mitochondrial DNA variation and population genetic structure of the blue crab *Callinectes sapidus* in the eastern United States. *Marine Biology* 145, 769–777.
- Melo G.A.S.** (1996) *Manual de identificação dos brachyura (caranguejos e siris) do litoral Brasileiro*. São Paulo: Plêiade/FAPESP, 604 pp.
- Mesquita A.R. and Harari J.** (1977) Variações sazonais em águas costeiras: Brasil Lat 24. Parte II. *Boletim do Instituto Oceanográfico de São Paulo* 26, 339–365.
- Ministério do Meio Ambiente.** (2004) Lista Nacional das espécies de invertebrados aquáticos e peixes sobreexploradas ou ameaçadas de sobreexploração. Instrução Normativa No. 5 de 21 de Maio de 2004. http://www.ibama.gov.br/sisbio/legislacao.php?id_arq=28.
- Ministério do Meio Ambiente.** (2005) *Boletim estatístico da pesca marítima e estuarina do nordeste do Brasil—2004. Centro de Pesquisa e Gestão de Recursos Pesqueiros do Litoral Nordeste—2004—Tamarandé, PE: CEPENE, 2004.* http://www.ibama.gov.br/cepene/download.php?id_download=316
- Miranda L.B.** (1970) Flutuações da Corrente do Brasil e variações da distribuição horizontal da temperatura na região costeira entre Cabo de São Tomé e Ilha de São Sebastião em Janeiro e Fevereiro e Abril de 1970. *Caderno de Ciências da Terra* 3, 13–14.
- Moritz C.** (1994) Defining 'evolutionarily significant units' for conservation. *Trends in Ecology and Evolution* 9, 373–375.
- Oliveira L.P.H.** (1946) Estudos ecológicos dos crustáceos comestíveis Uçá e Guaiamú, *Cardisoma guanhumi* Latreille e *Ucides cordatus* (L). Gecarcinidae, Brachyura. *Memórias do Instituto Oswaldo Cruz* 44, 395–322.
- Oliveira-Neto J.F., Boeger W.A., Pie M.R., Ostrensky A. and Hungria D.B.** (2007) Genetic structure of populations of the mangrove crab *Ucides cordatus* (Decapoda: Ocypodidae) at local and regional scales. *Hydrobiologia* 583, 69–76. doi 10.1007/s10750.
- Oliveira-Neto J.F., Pie M.R., Boeger W.A., Ostrensky A. and Baggio R.A.** (2008) Population genetics and evolutionary demography of *Ucides cordatus* (Decapoda: Ocypodidae). *Marine Ecology* 28, 1–10.
- Pfeiler E., Hurtado L.A., Knowles L.L., Torre-Cosío J., Bourillón-Moreno L., Márquez-Farías J.F. and Montemayor-López G.** (2005) Population genetics of the swimming crab *Callinectes bellicosus* (Brachyura: Portunidae) from the eastern Pacific Ocean. *Marine Biology* 146, 1432–1793.
- Posada D., Crandall K.A. and Templeton A.R.** (2000) GEODIS: a program for the cladistic nested analysis of the geographical distribution of genetic haplotypes. *Molecular Ecology* 9, 487–488.
- Rogers A.J. and Harpending H.** (1992) Population growth makes waves in the distribution of pairwise genetic differences. *Molecular Biology and Evolution* 9, 552–569.
- Roman J. and Palumbi S.R.** (2004) A global invader at home: population structure of the green crab, *Carcinus maenas*, in Europe. *Molecular Ecology* 13, 2891–2898. doi: 10.1111/j.1365-294X.2004.02255.x.
- Schneider S. and Excoffier L.** (1999) Estimation of demographic parameters from distribution of pairwise differences when the mutation rates vary among sites: application to human mitochondrial DNA. *Genetics* 152, 1072–1089.
- Seutin G., White B.N. and Boag P.T.** (1991) Preservation of avian blood and tissues samples for DNA analysis. *Canadian Journal of Zoology* 69, 82–90.
- Silva A.C., Araújo M. and Bourlès B.** (2005) Variação sazonal da estrutura de massas de água na plataforma continental do amazonas e área oceânica adjacente. *Revista Brasileira de Geofísica* 23, 145–157.
- Slatkin M. and Hudson R.R.** (1999) Pairwise comparisons of mitochondrial DNA sequences in stable and exponentially growing populations. *Genetics* 129, 555–562.
- Tajima F.** (1989) Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123, 585–595.
- Templeton A.R., Routman E. and Phillips C.A.** (1995) Separating population structure from population history: a cladistic analysis of the geographical distribution of mitochondrial DNA haplotypes in the tiger salamander, *Ambystoma tigrinum*. *Genetics* 140, 767–782.
- Weber L.I. and Levy J.A.** (2000) Genetic population structure of the swimming crab *Callinectes danae* (Crustacea: Decapoda) in southern Brazil. *Hydrobiologia* 420, 203–210.
- Weinberg J.R., Dahlgren T.G., Trowbridge N. and Halanych K.M.** (2003) Genetic differences within and between species of deep-sea crabs (*Chaceon*) from the north Atlantic ocean. *Biological Bulletin. Marine Biological Laboratory, Woods Hole* 204, 318–326.
- Weisstein E.W.** 'Bonferroni Correction.' In *MathWorld—A Wolfram Web Resource*. <http://mathworld.wolfram.com/BonferroniCorrection.html>
- and
- Wilson G.A. and Rannala B.** (2003) Bayesian inference of recent migration rates using multilocus genotypes. *Genetics* 163, 1177–1191.

Correspondence should be addressed to:

Marcio R. Pie
 Departamento de Zoologia, Caixa Postal 19073
 Universidade Federal do Paraná. Curitiba, PR 81531-990,
 Brazil
 email: pie@ufpr.br